

Stop search using boosted tops

Michihisa Takeuchi

Universität Heidelberg

at Beijing, 14th Aug 2012

Top at the LHC


closest to new physics \rightarrow probe for new physics

- fine tuning problem
 - \rightarrow cancellation via top partner
- Tevatron anomalies (A_{FB}^t , single top etc.)
- copiously produced via strong interaction at LHC

7,8TeV LHC $\sim 2,000,000 t\bar{t}$

Tevatron $\sim 40,000 t\bar{t}$


\rightarrow precision physics



Top at the LHC

closest to new physics \rightarrow probe for new physics

- fine tuning problem
 - \rightarrow cancellation via top partner
- Tevatron anomalies (A_{FB}^t , single top etc.)
- copiously produced via strong interaction at LHC
 - 7,8TeV LHC $\sim 2,000,000 \bar{t}t$
 - Tevatron $\sim 40,000 \bar{t}t$ \rightarrow precision physics



HEPTopTagger: hadronic top $t \rightarrow 3j$


- full momentum reconstruction possible in principle
 - \rightarrow important beyond discovery
- top against 10^3 larger QCD, how to identify?
 - $\sigma_{t\bar{t}}^{14\text{TeV}} = 918 \text{ pb} \leftrightarrow \sigma_{3j}^{14\text{TeV}} \sim 2 \cdot 10^6 \text{ pb}$
- take 3 jets with simple m_t, m_W condition
 - \rightarrow large QCD combinatorial BG kill us

Boosted Tops at the LHC

top jet


- top at rest \rightarrow separate 3 jets
- boosted top \rightarrow massive jet

$$R \sim 2m/p_T$$



QCD jet

- 2 jet events dominate QCD
- soft-collinear nature in its substructure
- take massive jet & look into jet substructure
combinatorics significantly reduced



Boosted Tops at the LHC

Boosted Top

HEPTopTagger


Applications

Summary

top as a probe

- new physics search with \cancel{E}_T
→ need recoil

- top at rest: not useful
- boosted tops: carry information on dark matter
better S/B (cf. M_{T2} end point.)



Boosted Tops at the LHC

Boosted Top

HEPTopTagger


Applications

Summary

top as a probe

- new physics search with \cancel{E}_T
→ need recoil

- top at rest: not useful
- boosted tops: carry information on dark matter
better S/B (cf. M_{T2} end point.)



top taggers

several top taggers available: focus on $p_T > 500$ GeV.

[Kaplan, Rehermann, Schwartz, Tweedie]


[Thaler, Wang]

[Almeida, Lee, Perez, Sterman, Sung]

Moderately Boosted Tops at the LHC

top p_T distribution


- $p_T > 500$ GeV: not many in SM
 $\sigma_{>200\text{GeV}} \sim 50\sigma_{>500\text{GeV}}$
- need top tagger valid down to
low p_T range \rightarrow testable



Moderately Boosted Tops at the LHC


top p_T distribution

- $p_T > 500$ GeV: not many in SM
 $\sigma_{>200\text{GeV}} \sim 50\sigma_{>500\text{GeV}}$
- need top tagger valid down to
low p_T range \rightarrow testable



- light top partners also provide
tops in the same range

we focus on $p_T > 200$ GeV
 \rightarrow need fat jet with $R = 1.5$



1. fat jets – C/A with $R = 1.5$, $p_T^{\text{fatjet}} > 200 \text{ GeV}$

Boosted Top

HEP**Top**Tagger



Applications

Summary


- fat jets** – C/A with $R = 1.5$, $p_T^{\text{fatjet}} > 200 \text{ GeV}$
- find subjets by mass drop criterion**

$$j = j_1 + j_2$$

$$m_j \gg m_{j_1}, m_{j_2} \text{ (decay)} \leftrightarrow m_j \sim m_{j_1} \gg m_{j_2} \text{ (QCD)}$$



- keep j_1 and j_2 for $m_{j_1} < 0.8m_j$ until $m_j < 50 \text{ GeV}$



HEPTopTagger [JHEP 1010:078,2010. arXiv:1006.2833 T. Plehn, M. Spannowsky, D. Zerwas, MT] [Phys.Rev. D85 (2012) 034029, arXiv:1111.5034]

Boosted Top

HEPTopTagger


Applications

Summary

1. **fat jets** – C/A with $R = 1.5$, $p_T^{\text{fatjet}} > 200$ GeV
2. **find subjects by mass drop criterion**
 - keep j_1 and j_2 for $m_{j_1} < 0.8m_j$ until $m_j < 50$ GeV
3. **take 3 subjects with best filtered mass**
 - $|m_{jjj}^{\text{filt}} - m_t| < 25$ GeV \rightarrow **top candidate**

filtering [Butterworth et al.]

- effect of pile-up, underlying events $\sim R^2$
- reduce effective area with smaller R_{filt} and n_{filt}




- $R_{\text{filt}} = \min\{0.3, R_{ij}/2\}$ and $n_{\text{filt}} = 5$ ($t \rightarrow bWg \rightarrow bgjjg$)

- 1. fat jets** – C/A with $R = 1.5$, $p_T^{\text{fatjet}} > 200$ GeV
- 2. find subjects by mass drop criterion**
 - keep j_1 and j_2 for $m_{j_1} < 0.8m_j$ until $m_j < 50$ GeV
- 3. take 3 subjects with best filtered mass**
 - $|m_{jjj}^{\text{filt}} - m_t| < 25$ GeV \rightarrow **top candidate**
- 4. check mass ratios**
 - 3 subjects: $p_1, p_2, p_3 \rightarrow m_{12}, m_{13}, m_{23}$
 - $m_t^2 = m_{123}^2 = m_{12}^2 + m_{13}^2 + m_{23}^2 \rightarrow$ 2D mass ratios

HEPTopTagger



[JHEP 1010:078,2010. arXiv:1006.2833 T. Plehn, M. Spannowsky, D. Zerwas, MT]
[Phys.Rev. D85 (2012) 034029, arXiv:1111.5034]

1. **fat jets** – C/A with $R=1.5$, $p_T^{\text{fatjet}} > 200$ GeV
2. **find subjects by mass drop criterion**
 - keep j_1 and j_2 for $m_{j_1} < 0.8m_j$ until $m_j < 50$ GeV
3. **take 3 subjects with best filtered mass**
 - $|m_{jjj}^{\text{filt}} - m_t| < 25$ GeV \rightarrow **top candidate**
4. **check mass ratios**
 - 3 subjects: $p_1, p_2, p_3 \rightarrow m_{12}, m_{13}, m_{23}$
 - $m_t^2 = m_{123}^2 = m_{12}^2 + m_{13}^2 + m_{23}^2 \rightarrow$ 2D mass ratios



- W mass condition, $\bar{t}\bar{t}$ soft-collinear cut \rightarrow **tagged top**
- no b -tag information



efficiency



- efficiency $\sim 30\%$ for hadronic tops, $2 \sim 4\%$ mis-tag rate
- momentum well reconstructed
- validation with ATLAS experimentalists in Heidelberg

[G. Kasieczka, S. Schätzel, A. Schöning]


Validation by ATLAS [ATLAS-CONF-2012-065]



- efficiency $\sim 30\%$ for hadronic tops, 2 $\sim 4\%$ mis-tag rate
- validation by ATLAS experimentalists in Heidelberg

[G. Kasieczka, S. Schätzel, A. Schöning]

- data well described by MC




Applications

Scalar Top Pairs at 14 TeV

hadronic mode [JHEP 1010 (2010) 078, T. Plehn, M. Spannowsky, MT, D. Zerwas]

- $\tilde{t}_1 \tilde{t}_1^* \rightarrow (t \tilde{\chi}_1^0)(\bar{t} \tilde{\chi}_1^0): m_{\tilde{t}_1} = 390 \text{ GeV}$
- main BG: $t\bar{t}$ +jets, W +jets and QCD



events in 1 fb^{-1}	$\tilde{t}_1 \tilde{t}_1^*$	$t\bar{t}$	QCD	W+jets	Z+jets	S/B	$S/\sqrt{B}_{10 \text{ fb}^{-1}}$
$m_{\tilde{t}} [\text{GeV}]$	390 440 490 540 640						390
$p_{T,j} > 200 \text{ GeV}, \ell \text{ veto}$	447 292 187 124 46	87850	$2.4 \cdot 10^7$	$1.6 \cdot 10^5$	n/a	$\sim 10^{-5}$	
$\cancel{E}_T > 150 \text{ GeV}$	234 184 133 93 35	2245	$2.4 \cdot 10^5$	1710	2240	$\sim 10^{-3}$	
first top tag	91 75 57 42 15	743	7590	90	114	0.01	
second top tag	12.4 11 8.4 6.3 2.3	32	129	5.7	1.4	0.07	
b -tag for 1 st top tag	7.4 6.3 5.0 3.8 1.4	19	2.6	$\lesssim 0.2$	$\lesssim 0.05$	0.34	5.0
$m_{T2} > 250 \text{ GeV}$	5.0 4.9 4.2 3.2 1.2	4.2	$\lesssim 0.6$	$\lesssim 0.1$	$\lesssim 0.03$	1.0	7.1

W+jets, Z+jets negligible with 2 top tag


QCD negligible with additional b -tag

$t\bar{t}$ reduced with m_{T2} cut

Scalar Top Pairs at 14 TeV


hadronic mode [JHEP 1010 (2010) 078, T. Plehn, M. Spannowsky, MT. D. Zerwas]

- $\tilde{t}_1 \tilde{t}_1^* \rightarrow (t \tilde{\chi}_1^0)(\bar{t} \tilde{\chi}_1^0): m_{\tilde{t}_1} = 390 \text{ GeV}$
- main BG: $t\bar{t}$ +jets, W +jets and QCD



events in 1 fb^{-1}	$\tilde{t}_1 \tilde{t}_1^*$	$t\bar{t}$	QCD	W+jets	Z+jets	S/B	S/\sqrt{B}	10 fb^{-1}
$m_{\tilde{t}}[\text{GeV}]$	390 440 490 540 640						390	
$p_{T,j} > 200 \text{ GeV}, \ell \text{ veto}$	447 292 187 124 46	87850	$2.4 \cdot 10^7$	$1.6 \cdot 10^5$	n/a	$\sim 10^{-5}$		
$\cancel{E}_T > 150 \text{ GeV}$	234 184 133 93 35	2245	$2.4 \cdot 10^5$	1710	2240	$\sim 10^{-3}$		
first top tag	91 75 57 42 15	743	7590	90	114	0.01		
second top tag	12.4 11 8.4 6.3 2.3	32	129	5.7	1.4	0.07		
b -tag for 1 st top tag	7.4 6.3 5.0 3.8 1.4	19	2.6	$\lesssim 0.2$	$\lesssim 0.05$	0.34		5.0
$m_{T2} > 250 \text{ GeV}$	5.0 4.9 4.2 3.2 1.2	4.2	$\lesssim 0.6$	$\lesssim 0.1$	$\lesssim 0.03$	1.0		7.1


W +jets, Z +jets negligible with 2
QCD negligible with additional ℓ
 $t\bar{t}$ reduced with m_{T2} cut



Scalar Top Pairs at 14 TeV

hadronic mode [JHEP 1010 (2010) 078, T. Plehn, M. Spannowsky, MT, D. Zerwas]

- $\tilde{t}_1 \tilde{t}_1^* \rightarrow (t \tilde{\chi}_1^0)(\bar{t} \tilde{\chi}_1^0): m_{\tilde{t}_1} = 390 \text{ GeV}$
- main BG: $t\bar{t}$ +jets, W +jets and QCD



events in 1 fb^{-1}	$\tilde{t}_1 \tilde{t}_1^*$	$t\bar{t}$	QCD	W+jets	Z+jets	S/B	S/\sqrt{B}	10 fb^{-1}
$m_{\tilde{t}} [\text{GeV}]$	390 440 490 540 640						390	
$p_{T,j} > 200 \text{ GeV}, \ell \text{ veto}$	447 292 187 124 46	87850	$2.4 \cdot 10^7$	$1.6 \cdot 10^5$	n/a	$\sim 10^{-5}$		
$\cancel{E}_T > 150 \text{ GeV}$	234 184 133 93 35	2245	$2.4 \cdot 10^5$	1710	2240	$\sim 10^{-3}$		
first top tag	91 75 57 42 15	743	7590	90	114	0.01		
second top tag	12.4 11 8.4 6.3 2.3	32	129	5.7	1.4	0.07		
b -tag for 1 st top tag	7.4 6.3 5.0 3.8 1.4	19	2.6	$\lesssim 0.2$	$\lesssim 0.05$	0.34		5.0
$m_{T2} > 250 \text{ GeV}$	5.0 4.9 4.2 3.2 1.2	4.2	$\lesssim 0.6$	$\lesssim 0.1$	$\lesssim 0.03$	1.0		7.1


$$S/B = 1, S/\sqrt{B} > 5 \text{ at } 14 \text{ TeV with } 10 \text{ fb}^{-1}$$

- stop mass from $m_{T2}(m_{\tilde{\chi}_1^0})$ endpoint [C. G. Lester, D. J. Summers]
like sleptons or sbottoms

Scalar Top Pairs at 14 TeV

hadronic mode [JHEP 1010 (2010) 078, T. Plehn, M. Spannowsky, MT, D. Zerwas]

- $\tilde{t}_1 \tilde{t}_1^* \rightarrow (t \tilde{\chi}_1^0)(\bar{t} \tilde{\chi}_1^0): m_{\tilde{t}_1} = 390 \text{ GeV}$
- main BG: $t\bar{t}$ +jets, W +jets and QCD



events in 1 fb^{-1}	$\tilde{t}_1 \tilde{t}_1^*$	$t\bar{t}$	QCD	W+jets	Z+jets	S/B	$S/\sqrt{B}_{10 \text{ fb}^{-1}}$
$m_{\tilde{t}} [\text{GeV}]$	390 440 490 540 640						390
$p_{T,j} > 200 \text{ GeV}, \ell \text{ veto}$	447 292 187 124 46	87850	$2.4 \cdot 10^7$	$1.6 \cdot 10^5$	n/a	$\sim 10^{-5}$	
$\cancel{E}_T > 150 \text{ GeV}$	234 184 133 93 35	2245	$2.4 \cdot 10^5$	1710	2240	$\sim 10^{-3}$	
first top tag	91 75 57 42 15	743	7590	90	114	0.01	
second top tag	12.4 11 8.4 6.3 2.3	32	129	5.7	1.4	0.07	
b -tag for 1 st top tag	7.4 6.3 5.0 3.8 1.4	19	2.6	$\lesssim 0.2$	$\lesssim 0.05$	0.34	5.0
$m_{T2} > 250 \text{ GeV}$	5.0 4.9 4.2 3.2 1.2	4.2	$\lesssim 0.6$	$\lesssim 0.1$	$\lesssim 0.03$	1.0	7.1

$$S/B = 1, S/\sqrt{B} > 5 \text{ at } 14 \text{ TeV with } 10 \text{ fb}^{-1}$$

- stop mass from $m_{T2}(m_{\tilde{\chi}_1^0})$ endpoint [C. G. Lester, D. J. Summers]
- like sleptons or sbottoms

semi-leptonic mode [JHEP 1105 (2011) 135 [arXiv:1102.0557], T. Plehn, M. Spannowsky, MT]

boosted leptonic top $S/B \sim 2, S/\sqrt{B} > 5 \text{ at } 14 \text{ TeV with } 10 \text{ fb}^{-1}$

Scalar Top Pairs at 8TeV

[arXiv:1205.2696 T. Plehn, M. Spannowsky, MT]

- $\sigma^{8\text{TeV}} \sim \frac{1}{10} \sigma^{14\text{TeV}}$: both for $t\bar{t}$ and $\tilde{t}_1\tilde{t}_1^*$
- 2 boosted tops: not enough signal left
- $t\bar{t}$: dominant background at the end

Boosted Top

HEPTopTagger


Applications

Summary

Scalar Top Pairs at 8TeV

[arXiv:1205.2696 T. Plehn, M. Spannowsky, MT]


- $\sigma^{8\text{TeV}} \sim \frac{1}{10} \sigma^{14\text{TeV}}$: both for $t\bar{t}$ and $\tilde{t}_1\tilde{t}_1^*$
- 2 boosted tops: not enough signal left
- $t\bar{t}$: dominant background at the end



Scalar Top Pairs at 8TeV

[arXiv:1205.2696 T. Plehn, M. Spannowsky, MT]

- $\sigma^{8\text{TeV}} \sim \frac{1}{10} \sigma^{14\text{TeV}}$: both for $t\bar{t}$ and $\tilde{t}_1\tilde{t}_1^*$
- 2 boosted tops: not enough signal left
- $t\bar{t}$: dominant background at the end



- 1 boosted top and 1 non-boosted top
 - hadronic mode: 1 hadronic top-tag + b -jet + \cancel{E}_T
 - semi-leptonic mode: 1 hadronic top-tag + ℓ, \cancel{E}_T


– semi-leptonic mode: 1 top-tag + ℓ, \cancel{E}_T

Boosted Top

HEPTopTagger

Applications

Summary



$$t\bar{t} \rightarrow t_h + b\ell\nu$$

negligible with $m_T(\ell, \cancel{E}_T) > 150$ GeV


$$t\bar{t} \rightarrow b\bar{b} + \tau_h\ell + 2\nu$$

fake hadronic top tag with ISR or τ_h

→ subjet id: b -tag, τ_h rejection.

$\sqrt{s} = 8$ TeV, $R = 1.5$	$\tilde{t}_1\tilde{t}_1^*$							$t\bar{t}$	$S/B S/\sqrt{B}_{10\text{fb}^{-1}}$	
m_T [GeV]	350	400	450	500	600	700		400		
cross section [fb]	760	337	160	80.5	23.0	7.19	$2.34 \cdot 10^5$			
$n_\ell = 1, \cancel{E}_T > 100$ GeV, $n_{\text{fat}} \geq 1$	104.37	61.49	34.81	19.54	6.28	2.11	5631			
$n_{\text{tag}} = 1$	13.09	9.02	5.80	3.60	1.33	0.50	788.79			
$m_T > 150$ GeV	4.63	4.27	3.25	2.19	0.94	0.38	3.28	1.0	6.5	
$j_b = b$	1.47	1.38	1.06	0.70	0.31	0.13	0.63	2.1	5.4	
$(j_b, j_{W1}, j_{W2}) = (b, j, j)$, reject τ_h	1.20	1.16	0.88	0.60	0.27	0.11	0.25	4.1	6.9	

– di-lepton mode



$$\bar{t}\bar{t} \rightarrow \bar{b}\bar{b} + \ell\ell + 2\nu$$

negligible with $m_{T2}^{\ell\ell} > 100$ GeV

$$m_{T2} = \min_{\cancel{E}_T \text{ split}} \left[\max \{ m_T^{\ell_1}, m_T^{\ell_2} \} \right]$$

$\sqrt{s} = 8$ TeV	$\tilde{t}_1\tilde{t}_1^*$						$\bar{t}\bar{t}$	$\bar{t}\bar{t}Z$	S/B	$S/\sqrt{B}_{10\text{fb}^{-1}}$
$m_{\tilde{t}} [\text{GeV}]$	350	400	450	500	600	700			400	
$n_\ell = 2$	30.98	14.27	7.07	3.58	1.04	0.33	7650.88	n.a.		
$\cancel{E}_T > 100\text{GeV}$	19.04	9.99	5.40	2.94	0.91	0.30	1312.74	0.35		
$m_{T2}^{\ell\ell} > 100$ GeV	6.05	4.30	2.70	1.65	0.56	0.20	0.65	0.09	5.8	16
$m_{T2}^{\ell\ell} > 150$ GeV	0.81	1.21	1.06	0.81	0.34	0.14	0.00	0.02	n.a.	n.a.

For scalar top mass 400 GeV for 10 fb^{-1}

- fully hadronic mode: statistically limited

$$S/B \sim 0.8, S/\sqrt{B} \sim 1.5 \text{ (two top tag)}$$

$$S/B \sim 1, S/\sqrt{B} \sim 3 \text{ (one top tag)}$$

- semi-leptonic mode:

$$S/B \sim 4, S/\sqrt{B} \sim 7$$

- di-lepton mode: not conclusive

$$S/B \sim 6, S/\sqrt{B} \sim 16$$

95% C.L. exclusion up to $\sim 600 \text{ GeV}$

Combining all channels important for discovery
possible in 2012 ! ($25 \sim 30 \text{ fb}^{-1}$)

Summary

HEPTopTagger available on <http://www.thphys.uni-heidelberg.de/~plehn/>

- moderate p_T tops ($> 200\text{GeV}$) \rightarrow testable in SM
- fat jets kill combinatorics
- jet substructure
 - thrown information \rightarrow use all available information
- momentum well reconstructed
- general idea: tops at LHC identified just like bottoms

Applications

- stop pairs at 14 TeV (2 boosted tops)
 $S/B \sim 1$ (hadronic), $S/B \sim 2$ (semi-leptonic), with $S/\sqrt{B} > 5$
- stop pairs at 8 TeV with 10fb^{-1}
 - $S/B \sim 1, S/\sqrt{B} \sim 1.5$ (hadronic)
 - $S/B \sim 4, S/\sqrt{B} \sim 7$ (semi-leptonic)
 - $S/B \sim 6, S/\sqrt{B} \sim 16$ (di-leptonic)
- Other applications
 - A_{FB}^t [Phys.Rev. D84 (2011) 054005, J. L. Hewett, J. Shelton, M. Spannowsky, T.M.P. Tait, MT]
 - single top σ_s vs. σ_t measurements [arXiv:1207.4787 F. Kling, T. Plehn, MT]